

FIRST RECORD OF DINOSAUR EGGSHELLS AND TEETH FROM
THE NORTH-WEST AFRICAN MAASTRICHTIAN (MOROCCO)

by

Géraldine GARCIA*, Rodolphe TABUCE**, Henri CAPPETTA**, Bernard MARANDAT**,
Ilhem BENTALEB**, Aziza BENABDALLAH*** & Monique VIANEY-LIAUD**

SUMMARY

	Page
Abstract, Résumé	60
Introduction	60
Geological setting	60
Systematics	61
Oofamily Megaoolithidae ZHAO, 1975	62
? Oofamily Laevisoolithidae MIKHAÏLOV, 1991	62
? Oofamily Prismatoolithidae HIRSCH, 1994	63
Oofamily Krokolithidae KOHRING & HIRSCH, 1996	64
Theropoda indet.	64
Discussion and conclusion	65
Acknowledgements	66
Bibliography	66
Plate 1	69

** Laboratoire de Géobiologie, Biochronologie et Paléontologie Humaine, UMR 6046, Université de Poitiers, 40 Avenue du Recteur Pineau, 86022 Poitiers cedex, France (Email : geraldine.garcia@univ-poitiers.fr)

** Institut des Sciences de l'Evolution, Université Montpellier II, cc 064, Place Eugène Bataillon, 34095 Montpellier cedex 05, France

*** Ministère du Commerce, de l'Energie et des Mines, 6208 Rabat Institut, Rabat, Maroc.

Mots-clés: coquilles d'oeufs, dents de théropodes, Maastrichtien, Maroc, Afrique.

Key-words: amniotic eggshells, theropod teeth, Maastrichtian, Morocco, Africa.

ABSTRACT

We report the discovery of amniotic eggshells and theropod teeth from the Late Cretaceous period in Morocco. The megaloolithid family represents the only known dinosaur egg remains in the north part of Africa and attests indirectly to the occurrence of sauropod dinosaurs (titanosaurids) in the Maastrichtian of Africa.

RESUME

La découverte de coquilles d'oeufs et de dents de théropodes est signalée dans le Crétacé terminal du Maroc. La famille des Megaloolithidae représente les seuls restes d'oeufs de dinosaures du Nord de l'Afrique et atteste indirectement de la présence de dinosaures sauropodes (titanosauridés) dans le Maastrichtien d'Afrique.

INTRODUCTION

Dinosaur eggs and eggshells are now reported in numerous Mesozoic gondwanan localities (Hirsch, 1989; Sahni *et al.*, 1994; Vianey-Liaud *et al.*, 1994; Vianey-Liaud *et al.*, 1997; Calvo *et al.*, 1997), but a few are known from Africa: in fact, the discoveries from this continent are very limited and located especially in the southern part and more particularly in the Lower Jurassic sediments of South Africa's Elliot Formation (Zelenitsky & Modesto, 2002). It is noticeably different for dinosaur bones which have been reported with a large distribution in different stratigraphic levels (Upper Triassic to Maastrichtian) in Africa: for example, from South Africa (Kennedy *et al.*, 1987; Durand, 2001), from Lesotho (Knoll, 2002), from Malawi (Jacobs *et al.*, 1990), from Niger (Greigert *et al.*, 1954; Sereno *et al.*, 1996; Sereno *et al.*, 1999), from Egypt (Rauhut & Werner, 1997)...

Moreover, from the Upper Cretaceous of Morocco, the youngest dinosaurian bones are Cenomanian (Lavocat, 1951; about 93 m.y.) and the Maastrichtian dinosaur history was not yet known, until our discoveries. Field researches in Central Morocco (Middle Atlas), initiated in 1999 by one of us (R. Tabuce) have resulted in the discovery of numerous micro-remains in several outcrops, that straddle the Maastrichtian/Paleocene transition. Among others, the dinosaur specimens include more than twenty well preserved megaloolithid eggshells and two theropod teeth. So, we relate the first notable record of the megaloolithid family in North Africa, which attests indirectly of the occurrence of sauropod dinosaurs (titanosaurids) in the Upper Cretaceous of Africa.

GEOLOGICAL SETTING

The eggshells occur in the Irbzer Formation exposed in the Oukdiksou syncline (Fig. 1). The lower part of this formation is a yellowish, cross-bedded, calcareous sandstone with indeterminate shark teeth. It is overlain by phosphatic marls with vertebrate bone-beds. The elasmobranch fauna is typically Maastrichtian based on *Cretolamna maroccana*, *Serratolamna serrata*, *Squalicorax pristodontus*, *Squalicorax bassanii*, *Ganopristis leptodon* and *Dalpiazia stromeri*. The occurrence of *S. bassanii*, present mainly in the lower to middle Maastrichtian deposits in the Ouled Abdoun and Gannour Phosphate basins of the Moroccan Meseta indicates an age older than the Upper Maastrichtian. Continental fauna and flora occur in light grey mudstones and marls conformably overlaying these strict marine deposits. Charophytes, associated with the eggshells, indicate a Maastrichtian or Early Paleocene age (Schudack & Herbib, 1995), but ostracodes (Colin com. pers.) and an indeterminate ammonite discovered near the Oudiksou syncline (Andreu & Tronchetti, 1996) provide a Maastrichtian age for the upper part of the Irbzer Formation.




Figure 1.— a: Map showing Morocco and the location of the fossiliferous sites (ACH1a and ACH2)-indicated by the black F- in the Middle Atlas Mountains; b: Section ACH2 of the South Oudiksou syncline with locations of the eggshells-bearing beds in the Irbzer Formation. In the Oudiksou syncline, the lithofacies and thickness of Cretaceous series vary strongly in time and space, even on short distances; it is clearly the case for ACH1a and ACH2.

SYSTEMATICS

Eggshells were either collected on exposures and/or by screen-washing of sediment samples. They have been prepared by ultrasonic cleaning, and some (as they are strongly encrusted) by acetic acid (5%) processing. Scanning electron microscopy was used to show the ultrastructure and characteristic features of the outer and inner surfaces of eggshell fragments. Microstructures were studied from thin petrographic sections. Measurements were taken on thin sections with a micrometer, and on fragments with calipers. The structural and parataxonomic terminology follows that of Mikhailov *et al.* (1996) and Mikhailov (1997). Studied specimens are catalogued as ACH- in the collection of the laboratory of Paleontology, University of Montpellier II. The structure of the eggshell described was directly compared with other eggshell material from southern France, Spain and India.

Oofamily MEGALLOOLITHIDAE ZHAO, 1975 (emend. 1979)
Oogenus MEGALLOOLITHUS VIANEY-LIAUD *et al.*, 1994

Megaloolithus maghrebiensis oosp. nov.

Material: 25 eggshell fragments.

Holotype: ACH2-05, single eggshell fragment (thin section and fragment) (Fig. 2d and e)

Type locality: Achloulj 2, Irbzer Formation, Middle Atlas, Morocco (Fig. 1)

Etymology: In reference to the location - the Maghreb (the French-speaking North Africa) - where the eggshells were found.

Diagnosis: *Megaloolithus* irregularly compactituberculate. Eggshell thickness from 1.7 to 2.5 mm. Elongation ratio (H/W) of the units from 2.5 to 3.7. Fan-shaped units with arched growth lines in the mammillae throughout the shell units. Units are well separated (node diameters 0.5 to 1.3 mm). In the areas where the nodes are fused in short chains and flat, the units are not clearly defined and the horizontal growth lines continue from one to the other. Pore canals are straight or linked by oblique bridges, as in *Megaloolithus siruguei*. Pore canal diameters between 75 to 130 μ m. Large eisospherite (0.15 to 0.3 mm) per node. Size and shape of the egg unknown.

Description and comparison

This oospecies has been found throughout the sampled Achloulj sections, at the same stratigraphical level in the upper five meters of the Irbzer Formation. They are found scattered in grey, sometimes reddish marls, generally associated with thin detritic intercalations.

The morphological characteristics (multicanalicate system pore, thickness range and compactituberculate pattern of the outer surface) are very similar to the European *Megaloolithus* species: *M. siruguei* (Garcia & Vianey-Liaud, 2001a). However it differs by the generally larger node diameters, the wider pore canals and more

pronounced fan-shaped units.

Defined narrower units, more important shell thickness and wider node diameter differentiate the new material from megaloolithid eggshell previously described from India [*Megaloolithus mohabeyi* (Vianey-Liaud *et al.*, 2003): 1.8 to 1.9 mm, H/W <3] and from Argentina (Chiappe *et al.*, 1998: 1 to 1.78 mm, H/W < 2).

A very small fragment from Achloulj 1a (Fig. 1) [thin section ACH1a-01] displaying a discretispherulitic microstructure, is referred to an undetermined species of *Megaloolithus*; It is thin (0.7 mm) with nodes separated by flat areas.

? Oofamily **LAEVISOOLITHIDAE** MIKHAILOV, 1991
Genus **SUBTILOLITHUS** MIKHAILOV, 1991

Tipoolithus achlouljensis oogen. nov., oosp. nov.

Material: 12 eggshell fragments.

Holotype: single eggshell fragment ACH2-06 (thin section and fragment) (Fig. 2a-c)

Type locality: Achloulj 2, Irbzer Formation, Middle Atlas, Morocco (Fig. 1)

Etymology: Tip in reference to the shape of nodes and after the name of the type locality: Achloulj.

Diagnosis: Eggshell thickness 0.40-0.65 mm, excluding the ornamentation; continuous layer: mammillary layer ratio = 2/1 to 1/1; ornamentation dispersituberculate, with high nodes (ratio node height / eggshell thickness = 0.37-0.45); node diameters 0.15-0.40 mm; pore openings situated from top of the nodes to their base, or between them; wide pore openings (90-160 μ m); angusticanalicate pore system.

Description and comparison

This oospecies resembles *Porituberoolithus warnerensis* (Zelenitsky *et al.*, 1996) from the Upper Cretaceous of Oldman Formation (Canada), in its general microstructure and thickness (0.5 to 0.65 mm), but the location of the pores is different, exclusively on the nodes. The nodes of *Tipoolithus* are sharp forming tips, generally isolated and rarely twinned or grouped by three, dispersed over the outer surface. On one fragment, the nodes seem to be aligned but the fragments are very small (less than 9 mm²). The pore system is angusticanalicate with probably very wide and nonbranching canals. *Tipoolithus* is very similar to *Subtiliolithus kachchhensis* (Khosla & Sahni, 1995) of the ornithoid-ratite eggshells from the Late Cretaceous of Anjar (Gujarat, India). This oospecies also exhibits dispersituberculate ornamentation, but is slightly thinner (0.34-0.45 mm) than *Tipoolithus* (0.4 to 0.65 mm) and its nodes are rounded and low.

? Oofamily **PRISMATOLITHIDAE** HIRSCH, 1994 emend.
? *PRISMATOLITHUS* ZHAO & LI, 1993

Description

Two very small fragments, less than 1 mm², from Achloulj 1A (ACH1a-02) studied in thin section, show a prismatic structure with units consisting of slender, interlocking prisms. They exhibit small circular nodes (0.15-0.17 mm in diameter) and an eggshell thickness varying from 0.29 to 0.35 mm. They could correspond to a theropod egglayer (Zelenitsky & Hills, 1996; Mateus *et al.*, 1997).

Oofamily **KROKOLITHIDAE** KOHRING & HIRSCH, 1996.

Description

One single fragment (about 2 mm²) from also Achloulj 1A (ACH1a-03) shows a typical crocodiloid microstructure. The wide shell units display irregular wedges and the characteristic horizontal layering. On the outer surface, the top of the units are flat and separated by wide and shallow valleys.

THEROPODA indet.

Material: two teeth ACH2-07 and ACH2-08 (Fig. 2 f-h)

Description

Only two fragmentary teeth have been found in the Achloulj 2 section (Fig. 1). The first is a fragment of an asymmetrical and laterally compressed tooth without serrations and slightly flattened on one side. The tooth tip is missing, as well as is base. It is 2.7 mm high, 1.6 mm long and 0.8 mm wide. The enamel is very thin (0.01 mm thick) and damaged by thin vertical cracks. The pulp cavity is small, less than 10% of the tooth. The second tooth shows the same shape, low ridges and thin vertical cracks. It differs only by the occurrence of denticles on its posterior edge.

These teeth closely resemble the morphotype "*Paronychodon*" (Carpenter, 1982; Sanz & Frances, 1992) by their morphology, their proportions and their measurements. The lateral profile is strongly asymmetrical, curving posteriorly. They are covered by coarse longitudinal ridges on the two sides of the crown. Three long and low ridges are observed on the flattest side of the crown which becomes faint to the base. On the most rounded side, only one smooth and shallow groove is hardly distinct. One specimen lacks serrations entirely but the other has some denticles on its posterior edge, as it can occur on *Paronychodon* teeth.

DISCUSSION AND CONCLUSION

The teeth were found in a transitional level within the lowest eggshell bearing level, close to Early Maastrichtian dated strata. They represent the first Late Cretaceous Theropoda remains found in continental African levels. They cannot be precisely referred to any theropod family. *Paronychodon* is assigned either to the Dromaeosauridae or to the Troodontidae (Currie *et al.*, 1990), but even the generic attribution is not quite certain. Nevertheless, both families are Laurasian and the occurrence of this taxon confirms communication during the Upper Cretaceous period between Africa and Laurasia. It was already suspected by the mixed character of the European reptile fauna with both Laurasian and Gondwanian origins (Vasse, 1995; Le Loeuff, 1993).

Records of African Mesozoic eggs are very limited, and before this report, no detailed accounts of North African Cretaceous eggshell have been published. We establish from our Moroccan locality the presence of at least 4 egg morphotypes, two of which may have been laid by theropods: the prismatoolithid shell is similar to that of eggshell referred to *Troodon* (Horner & Weishampel, 1996) and the new genus *Tipoolithus* which shows a "ratite" morphotype could belong to the Laevisoolithid family probably laid either by enantiornithine birds and/or small theropods (Mikhailov, 1997). Thus, indirectly one or two small carnivorous dinosaurs (Theropoda) or birds can be listed in the Late Cretaceous dinosaur fauna of Achloulj.

The megaloolithid family is well diversified and common in the Campano-Maastrichtian continental deposits of Gondwana: Indian, South Europe (Zhao, 1975; Sahni *et al.*, 1994; Vianey-Liaud *et al.*, 1994; Vianey-Liaud & Lopez-Martinez, 1997) and South American localities [Peru (Vianey-Liaud *et al.*, 1997), Argentina (Chiappe *et al.*, 1998; 2000) and Brasil (Price, 1951; Magalhaes Ribeiro, 2000)]. *Megaloolithus maghrebiensis* nov. sp. from Morocco shows close affinities, but also differences especially in some quantitative criteria (node diameter, thickness...) with the South-American, Indian and European Cretaceous megaloolithid oospecies, well studied and described in detail (see for example Garcia, 1998; Chiappe *et al.*, 1998; Garcia & Vianey-Liaud, 2001a; Vianey-Liaud *et al.*, 2003; Garcia *et al.*, submitted). The similarities in egg taxa suggest close phyletic relationships. The discovery in Argentina of embryo remains related to titanosaurs (sauropod) inside typical discritispherulitic eggs (Chiappe *et al.*, 1998) is strongly significant, suggesting that all megaloolithids are associated to titanosaur-like egglayers. Thus, our Moroccan megaloolithid material, which can be tentatively attributed to titanosaurid dinosaur, fills a geographical gap in Gondwanan Late Cretaceous sauropod distributions. Because these eggshells were found in levels overlying Lower Maastrichtian marine strata, it is possible to confirm the presence of titanosaurs in the North African Maastrichtian. This was long known in India, and subject to discussion in Europe, where the Maastrichtian bone localities are scarce (Le Loeuff, 1991), and where the megaloolithid species diversity drastically decreases at the beginning of this stage (Garcia & Vianey-Liaud, 2001b). This discovery is important in the knowledge of the evolution of dinosaur diversity just before their extinction. As several successive levels have yielded bone and eggshell fragments, we can expect to increase rapidly this new fossil record in this crucial time span, near the

Cretaceous-Tertiary boundary.

ACKNOWLEDGEMENTS

The members of the mission would thank Mr M. Sadiqui, the head of the geology department and Mr L. Tabit, manager of the "division SIG et patrimoine géologique" for their helpful collaboration. This work was funded by the UMR 5554, "Institut des Sciences de l'Evolution and by the ECLIPSE CNRS program. Contribution ISEM n° 2002-109.

BIBLIOGRAPHY

- ANDREU, B. & TRONCHETTI, G., 1996. — Ostracodes et Foraminifères du Crétacé supérieur du synclinal d'El Koubbat, Moyen-Atlas, Maroc. *Geobios*, 29: 45-71.
- CALVO, J.O., ENGELLAND, S., HEREDIA, S.E. & SALGADO, L., 1997. — First record of dinosaur eggshells (?Sauropoda-Megaloolithidae) from Neuquén, Patagonia, Argentina. *Gaia*, 14: 23-32.
- CARPENTER, K.; 1982. — Baby dinosaurs from the Late Cretaceous Lance and Hell Creek formations and a description of a new species of theropod. *Cont. Geology* University of Wyoming, 20 (2): 123-134.
- CHIAPPE, L.M., CORIA, R.A., DINGUS, L., JACKSON, F., CHINSAMY, A. & FOX, M., 1998. — Sauropod dinosaur embryos from the Late Cretaceous of Patagonia. *Nature*, 396: 258-261.
- CHIAPPE, L.M., DINGUS, L., JACKSON, F., GRELLET-TINNER, G., CORIA R., LOOPE, D., CLARKE, J. & GARRIDO, A., 2000. — Sauropod eggs and embryos from the Late Cretaceous of Patagonia. *In*: BRAVO, A. M. & REYES, T. (eds.): 1st International Symposium on Dinosaurs Eggs and Babies, Extended abstracts: 23-30. Isona i Conca Della, Catalonia, Spain.
- CURRIE, P.J., RIGBY, J.K. Jr. & SLOAN, R.E., 1990. — Theropod teeth from the Judith River Formation of southern Alberta, Canada. *In*: CARPENTER, K. & CURRIE, P. (eds.): Dinosaur systematics: 108-125. Cambridge university Press.
- DURAND, J.F., 2001. — The oldest juvenile dinosaurs from Africa. *Afr. Earth Sci.*, 33: 597-603.
- GARCIA, G., 1998. — Les coquilles d'œufs de dinosaures du Crétacé supérieur du Sud de la France: Diversité, paléobiologie, biochronologie et paléoenvironnements. Ph.D. Thesis, Univ. Montpellier II, France. Unpublished
- GARCIA, G. & VIANEY-LIAUD, M., 2001a. — Nouvelles données sur les coquilles d'œufs de dinosaures de Megaloolithidae du sud de la France: systématique et variabilité intraspécifique. *C. R. Acad. Sci. Paris*, 332: 185-191.
- GARCIA, G. & VIANEY-LIAUD, M., 2001b. — Dinosaur eggshells as new biochronological markers in Late Cretaceous continental deposits. *Palaeogeog., Palaeoclim., Palaeoecol.*, 169: 153-164.
- GARCIA, G., CODREA, V. VAN ITTERBEECK, J., GODEFROIT, P. & SMITH, T., (submitted). — Megaloolithid eggs from Romania. *Act. Paleont. Polonica*.
- GREIGERT, J., JOULIA, J. & LAPPARENT, A.F. de., 1954. — Répartition stratigraphique des gisements de Vertébrés dans le crétacé du Niger. *C. R. Acad. Sci. Paris*, 239: 433-435.
- HERBIG, H. G. & FECHNER, G. G., 1994. — Cretaceous and Early Tertiary stratigraphy, facies and

- palynology of the eastern Bou Angueur syncline, Middle Atlas Mountains, Morocco. *Z. Deutsch. Geol. Gesellschaft*, 145: 249-273.
- HIRSCH, K.F., 1989. — Interpretations of Cretaceous and pre-Cretaceous eggs and eggshell fragments. In: GILLETTE, D. & LOCKLEYS, M. (eds.): *Dinosaurs tracks and traces*: 89-87. Cambridge University Press, New York.
- HORNER, J.R. & WEISHAMPEL, D.B., 1996. — A comparative embryological study of two ornithischian dinosaurs. *Nature*, 383: 103.
- JACOBS, L.L., WINKLER, D.A., KAUFULU, Z.M. & DOWNS, W.R., 1990. — The dinosaur beds of Malawi, Africa. *National Geog. Res.*, 6: 196-204.
- KENNEDY, W.J., KLINGER, H.C. & MATEER, N.J., 1987. — First record of an Upper Cretaceous sauropod dinosaur from Zululand, South Africa. *S. Afr. J. Sci.*, 83: 173-174.
- KHOSLA, A. & SAHNI, A., 1995. — Parataxonomic classification of Late Cretaceous dinosaur eggshells from India. *J. Paleont. Soc. India*, 40: 87-102.
- KNOLL, F., 2002. — Nearly complete skull of *Lesothosaurus* (Dinosauria: Ornithischia) from the Upper Elliot Formation (Lower Jurassic: Hettangian) of Lesotho. *J. Vert. Pal.*, 22 (2): 238-243.
- LAVOCAT, R., 1951. — Découverte de restes d'un grand dinosaurien Sauropode dans le crétacé du sud Marocain. *C. R. Acad. Sci. Paris*, 232: 169-170.
- LE LOEUFF, J., 1991. — The Campano-Maastrichtian vertebrate faunas from Southern Europe and their relationships with other faunas in the world: paleobiogeographical implications. *Cret. Res.*, 12: 93-114.
- LE LOEUFF, J., 1993. — European Titanosaurids. *Rev. Paleobiologie*, 7: 105-117.
- RAUHUT, O.W.M. & WERNER, C., 1997. — First record of a Maastrichtian sauropod dinosaur from Egypt. *Paleont. Africana*, 34: 63-67.
- MAGALHAES RIBEIRO, C.M., 2000. — Microstructural analysis of dinosaur eggshells from Bauru basin (Late Cretaceous), Minas Gerais, Brasil. In: BRAVO, A. M. & REYES, T. (eds.): 1st International Symposium on Dinosaurs Eggs and Babies, Extended abstracts: 117-122. Isona i Conca Della, Catalonia, Spain.
- MATEUS, I., MATEUS, M., MATEUS, O., TAQUET, P., RIBEIRO, V. & MANUPELLA, G., 1997. — Couvées, oeufs et embryons d'un dinosaure Théropode du Jurassique supérieur de Lourinha (Portugal). *C. R. Acad. Sci. Paris*, 325: 71-78.
- MIKHAILOV, K.E., 1997. — Fossil and recent eggshell in amniotic vertebrates: fine structure, comparative morphology and classification. *Sp. Pap. Paleontology*: 5-80.
- MIKHAILOV, K.E., BRAY, E. & HIRSCH, K., 1996. — Parataxonomy of fossil egg remains (Veterovata): principles and application. *J. Vert. Pal.*, 16: 763-769.
- PRICE, L., 1951. — Un ovo de dinosauro na formacao Bauru do Cretacio de Estado de Minas Gerais. Departamento Nacional da producao Mineral, Divisao de Geologia e Minería, *Notas prelim. Estudos*, 53: 1-9.
- SAHNI, A., TANDON, S.K., JOLLY, A., BAJPAI, S., SOOD, A. & SRINIVASAN, S., 1994. — Upper Cretaceous dinosaur eggs and nesting sites from the Deccan-volcano sedimentary province of peninsular India. In: CARPENTER, K., HIRSCH, K. & HORNER J.R. (eds.), *Dinosaur Eggs and Babies*: 204-226. Cambridge University Press.
- SANZ, J.L. & FRANCES, V., 1992. — Reptiles and mammals from the Late cretaceous new locality Quintanilla del Coco (Burgos Province, Spain). *N. Jb. Geol. Paläont. Abhandlungen*, 184 (3): 279-314.
- SCHUDACK, M.E. & HERBIB, H.G., 1995. — Charophytes from the Cretaceous-Tertiary boundary beds of the Middle Atlas Mountains, Morocco. *Geol. Méditerranéenne*, 22: 125-139.

- SERENO, P.C., DUTHEIL, D.B., LAROCHE, M., LARSSON, H.C.E., LYON, G.H., MAGWENE, P.M., SIDOR, C.A., VARRICHIO, D.J. & WILSON, J.A., 1996.— Predatory Dinosaurs from the Sahara and Late Cretaceous faunal differentiation. *Science*, 272: 986
- SERENO, P.C., BECK, A.L., DUTHEIL, D.B., LARSSON, H.C.E., LYON, G.H., MOUSSA, B., SADLER, R.W., SIDOR, C.A., VARRICHIO, D.J., WILSON, G.P. & WILSON J.A., 1999. — Cretaceous sauropods from the Sahara and the uneven rate of skeletal evolution among dinosaurs. *Science*, 286: 1342-1347.
- VASSE D., 1995. — *Ischyrochampsia meridionalis* n.g. n. sp., un crocodilien d'affinité gondwanienne dans le Crétacé supérieur du sud de la France. *N. Jb. Geol. Paläont. Mh.*, 8: 501-512.
- VIANEY-LIAUD, M., MALLAN, P., BUSCAIL, O. & MONTGELARD, C., 1994. — Review of French dinosaur eggshells : morphology, structure, mineral, and organic composition. In: CARPENTER, K., HIRSCH, K. & HORNER J.R. (eds.), *Dinosaur Eggs and Babies*: 151-183. Cambridge University Press.
- VIANEY-LIAUD, M., HIRSCH, K.F., SAHNI, A. & SIGE, B., 1997. — Late Cretaceous Peruvian eggshells and their relationships with Laurasian and eastern Gondwanan material. *Geobios*, 30 (1): 75-90.
- VIANEY-LIAUD, M. & LOPEZ-MARTINEZ, N., 1997. — Late Cretaceous dinosaur eggshells from the Tremp Basin, southern Pyrenees, Lleida, Spain. *J. Paleontol.*, 71 (6): 1157-1171.
- VIANEY-LIAUD, M., KHOSLA, A. & GARCIA, G., 2003. — Comparisons of European and Indian dinosaurs eggshells: paleobiogeographical implications. *J. Vert. Pal.*, 23 (3): 575-585
- ZELENITSKY, D.K. & HILLS L.V., 1996. — An egg clutch of *Prismatoolithus levis* oosp. nov. from the Oldman formation (Upper Cretaceous), Devil's Coulee, Southern Alberta. *Can. J. Earth Sci.*, 33: 1655-1667.
- ZELENITSKY, D.K., HILLS, L.V. & CURRIE, P.J., 1996. — Parataxonomic classification of ornithoid eggshell fragments from the Oldman Formation (Judith River Group; Upper Cretaceous), southern Alberta. *Can. J. Earth Sci.*, 33: 1655-1667.
- ZELENITSKY, D. K. & MODESTO, S.P., 2002. — Re-evaluation of the eggshell structure of eggs containing dinosaur embryos from the Lower Jurassic of South Africa. *S. Afr. J. Sci.*, 98: 407-408.
- ZHAO, Z., 1975. — The microstructure of the dinosaur eggshells of Nanshiung, Kwantung. *Vert. Palasiatica*, 13: 105-117.

PLATE 1

Tipoolithus achloujensis nov. oogen. nov. oosp. a. Drawing of thin section, showing the two layers (mammillary layer, with spherulites, and the continuous spongy layer) Scale = 1 mm. b. detail of a, PLM, natural light, showing the growth lines continuous in the "spongy" layer, the mammillary layer, and the wide pore canal; Scale = 0.5 mm. c: SEM of outer surface, a pit and a pore in a flat area between the nodes; Scale = 0.1mm.

Megaloolithus maghrebiensis nov. oogen. nov. oosp. d: Drawing of the holotype (thin section). e: Detail of a thin section of *M. maghrebiensis*, PLM, natural light, showing the regular arched growth lines, dipping at the pore canal level, the large eisospherites and circular growth lines around them and the pore canal. Scale 1 mm.

Teeth of Theropoda. f: first tooth, most flat side, with three faint ridges; g: anterior side, showing the faint edge, worn at the top; h: second tooth, convex side with denticles on the posterior edge. Scale = 1 mm.

